



THE EFFECT OF MAGNESIUM IN CHEMICAL EQUILIBRIUM FORMS OF POTASSIUM AND THE GROWTH OF YELLOW CORN IN TOW SOILS OF DIFFERENT TEXTURE

Mansour F. M. Al - Kilaby ¹, Abdul Baqi D.S. Al Maamouri ²

¹Assistant Lecturer, Department of Soil and Water Sciences, College of Agriculture, University of Kufa, Iraq, mansourf.alkilaby@uokufa.edu.iq

²Assistant Professor PhD., Department of Soil Sciences and Water Resources, College of Agricultural Engineering Sciences, University of Baghdad, Iraq abdulbaqi@coagri.uobaghdad.edu.iq

Received 21/ 11/ 2023, Accepted 31/ 1/ 2024, Published 31/ 12/ 2025



This work is licensed under a CCBY 4.0 <https://creativecommons.org/licenses/by/4.0>

ABSTRACT

This study was conducted to determine the effect of adding magnesium on the chemical equilibrium of potassium. The study included five levels of magnesium, Mg₀, Mg₁, Mg₂, Mg₃, and Mg₄, and two levels of potassium, K₀ and K₁₀₀, in two soils of different textures. The results of the study showed that adding magnesium contributed to increasing soluble, exchangeable, and non-exchangeable potassium. And by 64.706, 105.882, 164.706, and 235.294% / 13.986, 29.680, 47.945, and 77.169% / 4.538, 8.372, 13.224, and 26.682% for the level Mg₁, Mg₂, Mg₃, and Mg₄ compared to Mg₀, respectively. The soil with a silt clay texture outperformed the soil Silt Clay Leom texture with percentages of 131.818, 15.074 and 8.254%, for dissolved, exchangeable and non-exchangeable potassium, respectively.

key words: dispersion Clay, release of potassium, exchanged potassium, Unexchanged potassium.

تأثير المغسيسيوم في صور الازان الكيميائي للبوتاسيوم ونمو الذرة الصفراء في تربتي ذات نسجة مختلفة

منصور فرحان مخلول الكلابي ¹ ، عبد الباقي داود سلمان المعموري ²

¹مدرس مساعد، قسم علوم التربة والمياه، كلية الزراعة، جامعة الكوفة، العراق، mansourf.alkilaby@uokufa.edu.iq

² الاستاذ المساعد الدكتور، قسم علوم التربة والموارد المائية، كلية علوم الهندسة الزراعية، جامعة بغداد، العراق، abdulbaqi@coagri.uobaghdad.edu.iq

الخلاصة

جريت هذه الدراسة لمعرفة تأثير اضافة المغسيسيوم في صور الازان الكيميائي لعنصر البوتاسيوم تضمنت الدراسة اضافة خمس مستويات من المغسيسيوم Mg₀ و Mg₁ و Mg₂ و Mg₃ و Mg₄ ومستويين من البوتاسيوم K₀ و K₁₀₀ في تربتين مختلفتي النسجة ، اظهرت نتائج الدراسة ان اضافة المغسيسيوم قد ساهم في زيادة البوتاسيوم الذائب والمتبادل وغير المتبادل وبنسبة 64.706 و 105.882 و 164.706 و 235.294 و 13.986 % و 29.680 و 47.945 و 77.169 و 4.538 و 8.372 و 13.224 و 26.682 % وللمستوى Mg₁ و Mg₂ و Mg₃ و Mg₄ بالمقارنة مع Mg₀ على التوالي وتفوقت التربة ذات النسجة طينية غرينية على التربة ذات النسجة مزيجية طينية غرينية وبنسبة 131.818 و 15.074 و 8.254% و للبوتاسيوم الذائب والمتبادل وغير المتبادل على التوالي .

الكلمات المفتاحية : تشتت ، تحرر البوتاسيوم ، البوتاسيوم المتبادل ، البوتاسيوم غير المتبادل



INTRODUCTION

Potassium is an essential nutrient in plants. Therefore, it has become necessary to know the condition and fate of the potassium added to the soil, the extent of the crops' response to it, and whether the added potassium, which may be subject to fixation in the soil, can be released again, how quickly it is released, and what are the factors affecting the speed of its release. The results also showed that 28-76% of the added potassium is subject to fixation in minerals clay. (Al-Shamare & Essa, 2020; AL-Taee & Al-Maamouri, 2020; AlKhalil & Essa, 2020; AlKhalil & Essa, 2021; Al-Shamary *et al.*, 2022). The chemical behavior of any element affects the effectiveness and behavior of some other elements, either directly through the process of competition for the surface of the complex exchange and displacement, or indirectly through influencing other soil properties Magnesium is one of the elements that has the ability to disperse soil particles, and the main reason for this specific effect is due to the large size of its hydrosphere compared to calcium and the small size of the real ion, as it is (0.078 nm) while calcium (0.106 nm), in addition to its role in influencing the process of Ion exchange, as it carries a dual charge, (Al-Marsoumi & Jarallah, 2019). Also, magnesium is mobile in the soil, unlike other cations, and because of the large volume of the hydrosphere, this results in a relatively high abundance of this element in the soil solution and thus a decrease in its deposition. Also, the mobility of magnesium in the soil and the extent of its availability to plants is determined not only by its quantity but also by soil characteristics When the pH Soil. When the soil is less than 6.0, it is exposed to leaching in acidic soils, cation exchange capacity, concentration of other ions, humidity, temperature, and agricultural management practices such as fertilization. The magnesium ion is characterized by behavior and physicochemical properties that distinguish it from the rest of the other ions existing in the soil, such as the ion charge, diffusion coefficient, ionic diameter, polarity, and its role in dispersion clay and its reflection on the equilibrium forms of potassium in the soil, its relationship to plant growth (Nafawa, 2017). The current study aims to study the effect of magnesium and texture on the chemical equilibrium of potassium and the growth of maize plants.

MATERIALS AND METHODS

A biological experiment was carried out on two soils of different textures (silt clay and silt clay Leom) after the soil was brought from the field, dried and ground with a sieve with holes diameter of 4 mm. Chemical and physical analyzes of the soil were also conducted according to what was stated in (Black, 1965, and Jackson, 1958), the characteristics of which are shown in Table 1. Magnesium was added at five levels in the form of magnesium sulphate (100, 75, 50, 25.0) kg.ha^{-1} . Potassium was also added at two levels (0 and 100) kg.ha^{-1} in the form of potassium sulphate (41% K) and 200% was added kg ha^{-1} of nitrogen in the form of urea (46% N) and 100 kg ha^{-1} of phosphorus in the form of superphosphate fertilizer (21 P%). The seeds of yellow corn, a local variety, were planted in plastic pots with a capacity of 25 kg of soil, at a rate of 5 seeds per pot. They thinned out after a week of Germination into three plants. The forms of potassium (soluble, exchangeable and non-exchangeable) were estimated 90 days after germination. The absorbed potassium, plant height and leaf area were also estimated.



Table (1): Some chemical and physical properties of soil

Traits	Measurement value	value (S ₁)	value (S ₂)
Ece	dS.m ⁻¹	3.80	2.04
pH		7.82	8.02
Ca ²⁺		9.88	5.25
Mg ²⁺		6.65	3.71
Na ⁺		4.60	2.30
K ⁺		0.34	0.18
Cl ⁻		7.30	4.73
HCO ₃ ⁻		5.00	3.51
SO ₄ ²⁻		12.85	6.08
CO ₃ ²⁻		NiL	NiL
CEC	Cmmol.Kg ⁻¹	35.40	28.20
Organic matter		7.12	6.30
Total Carbonate		265.8	278.6
Active Carbonate		175	100
Clay		423.00	374.90
Silt		462.00	524.50
Sand		115.00	100.60
Texture		Silt Clay	Silt Clay Leom
Soluble K ⁺	Cmmol.Kg ⁻¹ Soil	0.03	0.02
Exchangeable K ⁺		0.27	0.24
Unexchangeable K ⁺		1.46	1.33
Mineral K ⁺		30.98	27.21
Total K ⁺		32.75	28.80

RESULTS AND DISCUSSION

1. The effect of soil texture and the levels of magnesium and added potassium in the concentration of soluble potassium cmmol K⁺.kg⁻¹ soil

Table (2) showed the effect of soil texture and the levels of magnesium and added potassium in the amount of soluble potassium. The results showed that there were significant differences between the treatments in increasing the amount of soluble potassium by 131.818% in the S₁ treatment compared to the S₂ treatment, and this was confirmed by (**Hamid et al., 2021**). On the role of soil texture, the amount of clay and silt, and its mineral composition in the amount of dissolved potassium in the soil, where there is a positive significant relationship with clay and silt and a negative relationship with sand in increasing the amount dissolved potassium in the soil. As for the effect of added magnesium levels, the results showed a significant increase in the amount of dissolved potassium by 64.706, 105.882, 164.706 and 235.294% for the levels Mg₁, Mg₂, Mg₃ and Mg₄ compared to the control treatment Mg₀, respectively. The results also showed that Mg₄ was superior to all levels of Mg in both soils. Which may have contributed to increasing the readiness of potassium through the role of magnesium in displacing potassium on the surface of soil particles and thus increasing its dissolved content due to concentration and equivalence. The results also showed that adding



potassium fertilizer at different levels led to a significant increase in soluble potassium by 165% in the K_{100} treatment compared to the K_0 control treatment, and this is consistent with what was indicated by (Al-Aqrabi, 2018; AlObaidi & Abdel-Rida, 2021). it Increasing the levels of potassium fertilizer added led to an increase in the amount of dissolved potassium. As for the bilateral interaction between soil texture and magnesium levels, the results showed significant differences between the different treatments, and the highest value in the S_1Mg_4 treatment was 0.071 cmmol $K^+ \cdot kg^{-1}$ soil and the lowest value when treated with S_2Mg_0 was 0.008 cmmol $K^+ \cdot kg^{-1}$ soil. This is due to the interplaying effect of soil texture and magnesium levels in the amount of dissolved potassium. The same trend also occurred with the bilateral interaction between the levels of added magnesium and potassium. The results showed significant differences between the different treatments, and the highest value in the $K_{100}Mg_4$ treatment was 0.082 cmmol $K^+ \cdot kg^{-1}$ soil and the lowest value when treated with K_0Mg_0 was 0.008 cmmol $K^+ \cdot kg^{-1}$ soil. The reason for this is attributed to the interfering effect of the levels of magnesium and potassium fertilizer in the amount of dissolved potassium. The bilateral interaction between soil texture and potassium levels had a significant effect in the amount of dissolved potassium, and the highest value when treated with S_1K_{100} was 0.075 cmmol $K^+ \cdot kg^{-1}$ soil and the lowest value when treated with S_2K_0 was 0.013 cmmol $K^+ \cdot kg^{-1}$ soil.

Table (2): Effect of soil texture and levels of magnesium and added potassium in the concentration of soluble potassium Cmmol $K^+ \cdot kg^{-1}$ soil.

S*K	Mg ₄	Mg ₃	Mg ₂	Mg ₁	Mg ₀	K	S
0.08	0.10	0.09	0.08	0.07	0.04	K100	S ₁
0.03	0.04	0.03	0.03	0.02	0.01	K0	
0.03	0.06	0.04	0.02	0.02	0.01	K100	S ₂
0.01	0.02	0.015	0.013	0.01	0.006	K0	
0.006	0.01					LSD	
S Average							
0.05	0.07	0.06	0.05	0.04	0.03	S1	S*Mg
0.02	0.04	0.03	0.018	0.01	0.01	S2	
0.004	0.004					LSD	
K Average							



0.05	0.08	0.07	0.05	0.04	0.03	K100	K*Mg
0.02	0.03	0.03	0.02	0.02	0.01	K0	
0.004	0.004					LSD	
	0.06	0.05	0.04	0.03	0.02	Mg Average	
	0.003					LSD	

As for the triple interaction between soil texture, magnesium levels, and potassium fertilizer, the results indicate that there are significant differences in the amount of dissolved potassium. The highest value appeared in the S₁K₁₀₀Mg₄ treatment, amounting to 0.101 cmmol K⁺. kg⁻¹ soil, while the lowest rate of soluble potassium in the S₂K₀Mg₀ treatment was 0.006 cmmol K⁺. kg⁻¹ soil.

2. Effect of soil texture and levels of magnesium and added potassium in the concentration of exchangeable potassium centimol K⁺.kg⁻¹ soil.

Tables (3) showed the effect of soil texture, magnesium fertilizer, and potassium fertilizer in the amount of potassium exchanged. The results showed that there were significant differences between the treatments. The results also indicated that the S₁ treatment was superior to S₂ due to its high clay content, which contributes to increasing the amount of potassium exchanged over The surface, as the silty clay texture outperformed the soil with a mixed clay-silty texture, by 15.074%, and this was confirmed by (Hamid *et al.*, 2021) about the role of soil texture and the amount of clay and silt in the amount of potassium exchanged in the soil, as there is a positive significant relationship with clay and silt. And negative with sand in increasing the amount of potassium exchanged in the soil. As for the effect of added magnesium levels, the results showed a significant increase in the amount of exchangeable potassium, as the percentage increase in exchangeable potassium for magnesium fertilizer levels reached 13.698, 29.680, 47.945, and 77.169% for the levels Mg₁, Mg₂, Mg₃, and Mg₄ compared to the control treatment Mg₀, respectively, which may have contributed In increasing the readiness of potassium through the role of magnesium in displacing potassium on the surface of soil particles and thus increasing its exchange rate due to concentration and equivalence. The results also showed that adding potassium fertilizer at different levels led to a significant increase in potassium exchanged in the K₁₀₀ treatment by 42.149% compared to the K₀ control treatment, and this is consistent with what was indicated by (Al-Aqrabi, 2018; Al-Obaidi & Abdel-Rida, 2021) on the role of added potassium in preparing exchangeable potassium during the growth stage, which led to an increase in the amount of exchangeable potassium. As for the bilateral interaction between the levels of soil texture and magnesium, the results showed significant differences between the different treatments, and the highest value in the S₁Mg₄ treatment was 0.436 cmmol K⁺.kg⁻¹ soil and the lowest value when treated with S₂Mg₀ was 0.205 cmmol K⁺.kg⁻¹ soil, the reason for this is attributed to the interplaying effect of soil texture and magnesium levels in the amount of potassium exchanged. The same trend also occurred with the bilateral interaction between the levels of added magnesium and potassium. The results showed significant differences between the different treatments, and the highest value in the K₁₀₀Mg₄ treatment was 0.436 cmmol K⁺.kg⁻¹ soil and the lowest value



when treated with K_0Mg_0 was $0.169 \text{ cmmol K}^+ \cdot \text{kg}^{-1}$ soil, the reason for this is attributed to the interfering effect of magnesium levels and potassium fertilizer in the amount of potassium exchanged. The bilateral interaction between soil texture and potassium levels had an effect Significant in the amount of potassium exchanged, the highest value when treated with S_1K_{100} was $0.362 \text{ cmmol K}^+ \cdot \text{kg}^{-1}$ soil and the lowest value when treated with S_2K_0 was $0.219 \text{ cmmol K}^+ \cdot \text{kg}^{-1}$ soil. As for the triple interaction, the results showed that there were significant differences between the treatments in the amount of potassium exchanged, as the highest rate of potassium exchanged in the $S_1K_{100} Mg_4$ treatment was $0.482 \text{ cmmol K}^+ \cdot \text{kg}^{-1}$ soil, while the lowest rate of exchangeable potassium in the $S_2K_0Mg_0$ treatment was $0.157 \text{ cmmol K}^+ \cdot \text{kg}^{-1}$ soil.

Table (3): Effect of soil texture and levels of added magnesium and potassium in the exchangeable potassium concentration $\text{Cmmol K}^+ \cdot \text{kg}^{-1}$ soil.

S*K	Mg ₄	Mg ₃	Mg ₂	Mg ₁	Mg ₀	K	S	
0.36	0.48	0.41	0.33	0.30	0.29	K_{100}	S_1	
0.26	0.39	0.29	0.25	0.21	0.18			
0.33	0.39	0.36	0.34	0.29	0.25	K_{100}	S_2	
0.22	0.29	0.24	0.21	0.20	0.16			
0.006	0.014					LSD		
S Average								
0.31	0.44	0.35	0.29	0.26	0.23	S_1	S^*Mg	
0.27	0.34	0.30	0.28	0.24	0.21			
0.005	0.01					LSD		
K Average								
0.34	0.44	0.39	0.33	0.29	0.27	K_{100}	K^*Mg	
0.24	0.34	0.26	0.23	0.21	0.17			
0.005	0.01					LSD		
	0.39	0.32	0.28	0.25	0.22	Mg Average		
	0.007					LSD		



3. The effect of soil texture and the levels of magnesium and added potassium in the concentration of unexchanged potassium centimol $K^+ \cdot kg^{-1}$ soil.

Tables (4) showed the effect of soil texture, magnesium fertilizer, and potassium fertilizer in the amount of non-exchangeable potassium. The results showed that there were significant differences between the treatments. The results also indicated that the S_1 treatment was superior to S_2 due to its high content of clay and non-exchangeable potassium, as the percentage of increase in The amount of non-exchangeable potassium in the S_1 treatment was 8.254% compared to the S_2 treatment. This was confirmed by (Hamid *et al.*, 2021) about the role of the amount of clay in increasing the amount of non-exchangeable potassium through the stabilization process between the layers.in addition to the type of clay mineral in the soil, which contribute It greatly increases the amount extracted from it in the analysis process. As for the effect of added magnesium levels, the results showed a significant increase in the amount of unexchanged potassium, as the percentage increase in unexchanged potassium reached 4.538, 8.372, 13.224, and 26.682% for the levels Mg_1 , Mg_2 , Mg_3 , and Mg_4 compared to the control treatment Mg_0 , respectively. This is attributed to the effect of magnesium on Dispersion of layers of minerals and the release of a portion of the potassium fixed between those layers. This is what was confirmed by (Nafawa ,2017) in the role of magnesium in dispersion soil particles due to the expansion of the volume of its hydrosphere and to the small crystal radius, which is 0.78A° . The results also showed that adding potassium fertilizer at different levels led to a significant increase in non-exchangeable potassium by 18.056% in the K_{100} treatment compared to the K_0 control treatment. This is consistent with what (Mahmood *et al.*, 2019) indicated about the role of the effect of the potassium fertilizer fraction added. In occupying the fixation sites, which leads to an increase in the amount of non-exchangeable potassium, and the excess is ready for absorption by the plant.

Table (4): Effect of soil texture and levels of added magnesium and potassium in concentration Unexchanged potassium Cmmol $K^+ \cdot kg^{-1}$ soil.

S^*K	Mg_4	Mg_3	Mg_2	Mg_1	Mg_0	K	S	
1.61	1.79	1.61	1.57	1.54	1.52	K100	S_1	
1.33	1.54	1.38	1.32	1.28	1.15	K0		
1.45	1.64	1.49	1.44	1.36	1.35	K100	S_2	
1.26	1.51	1.32	1.21	1.16	1.10	K0		
0.006	0.014					LSD		
S Average								
1.47	1.66	1.49	1.45	1.41	1.33	S1	S^*Mg	
1.36	1.57	1.40	1.32	1.26	1.22	S2		
0.004	0.010					LSD		



K Average							
1.53	1.71	1.55	1.50	1.45	1.43	K100	K*Mg
1.30	1.52	1.35	1.27	1.22	1.12	K0	
0.004	0.010						LSD
	1.62	1.45	1.39	1.34	1.28	Mg Average	
	0.007		LSD				

As for the bilateral interaction between soil texture and magnesium levels, the results showed significant differences between the different treatments, and the highest value in the S_1Mg_4 treatment was 1.664 cmmol $K^+ \cdot kg^{-1}$ soil and the lowest value when treated with S_2Mg_0 was 1.224 cmmol $K^+ \cdot kg^{-1}$ soil. This is due to the interplaying effect of soil texture and magnesium levels in the amount of non-exchangeable potassium. The same trend also occurred with the bilateral interaction between the levels of added magnesium and potassium. The results showed significant differences between the different treatments, and the highest value in the $K_{100}Mg_4$ treatment was 1.714 centimoles $K^+ \cdot Kg^{-1}$ soil and the lowest value when treated with K_0Mg_0 was 1.124 cmmol $K^+ \cdot kg^{-1}$ soil, the reason for this is attributed to the interfering effect of magnesium levels and potassium fertilizer in the amount of unexchanged potassium. The bilateral interaction between soil texture and potassium levels had a significant effect in the amount of unexchanged potassium, and the highest value when treated with S_1K_{100} was 1.606 cmmol $K^+ \cdot kg^{-1}$ soil and the lowest value when treated with S_2K_0 was 1.259 cmmol $K^+ \cdot kg^{-1}$ soil. As for the triple intervention, the results showed that there were significant differences between the treatments in the amount of non-exchanged potassium, as the highest rate of non-exchanged potassium in the $S_1K_{100}Mg_4$ treatment reached 1.787 centimol $K^+ \cdot kg^{-1}$ soil, while the lowest rate of non-exchangeable potassium in the $S_2K_0Mg_0$ treatment was 1.101 cmmol $K^+ \cdot kg^{-1}$ soil.

4. The effect of soil texture and levels of magnesium and added potassium in the amount of potassium absorbed, mg K.plant⁻¹

Figure (1) shows the effect of soil texture, magnesium fertilizer, and potassium fertilizer in the amount of potassium absorbed (mg K. plant⁻¹). The results showed that the S_1 treatment was superior to S_2 due to its higher content of clay and ready-made potassium compared to S_2 , as the percentage of increase in the amount of absorbed potassium was In the S_1 treatment, 11.111% respectively compared to the S_2 treatment, and this was confirmed by (Hamid *et al.*, 2021) . about the role of soil texture and the amount of clay in the properties of the soil and its ability to supply plants With nutrients and its effect on increasing the amount of potassium absorbed. As for the effect of magnesium fertilizer levels, the results showed that there were significant differences in the amount of potassium absorbed, as the percentage increase in the amount of potassium absorbed when using magnesium fertilizer levels reached 5.488, 12.670, 21.815, and 39.597% for the levels Mg_1 , Mg_2 , Mg_3 , and Mg_4 compared to the comparison treatment Mg_0 on respectively, this is consistent with what was obtained by (Al-Fadhy *et al.*, 2019) who showed that adding levels of magnesium positively affected the amount of potassium absorbed in the plant. The results also showed that there were significant

differences between the treatments. Adding potassium fertilizer at different levels significantly affected the amount of potassium absorbed by the plant, as the percentage increase in the amount of potassium absorbed in the K_{100} treatment when using potassium fertilizer levels was 20.262% compared to the comparison treatment K_0 , and the reason for this is Added potassium increases the efficiency of the photosynthesis process, then increases the weight of the plant and absorbs the largest amount of nutrients during the growth stage. This is consistent with what was indicated by (Asaad & Abdel-Rasoul, 2017; Elfahdawi *et al.*, 2020). As for the bilateral interaction between soil texture and magnesium levels, the results showed significant differences between the different treatments. The reason for this is attributed to the overlapping effect of soil texture and magnesium levels in the amount of potassium absorbed. The same trend also occurred when the bilateral interaction between magnesium levels and potassium fertilizer in the soil. The results showed significant differences. between different transactions this is due to the interfering effect of magnesium and potassium fertilizer levels in the amount of potassium absorbed. As for the bilateral interaction between soil texture and potassium levels, the results showed significant differences between the different treatments. The reason for this is attributed to the overlapping effect of soil texture and potassium levels in the amount of potassium absorbed. As for the three-way interaction between soil texture and the levels of added magnesium and potassium, the results indicate that there are significant differences in the amount of potassium absorbed, as the highest rate in the amount of potassium absorbed in the $S_1 K_{100} Mg_4$ treatment reached $7533.51 \text{ mg K.plant}^{-1}$, while the lowest rate was in the amount The potassium absorbed in the $S_2 K_0 Mg_0$ treatment amounted to $3987.92 \text{ mg K. plant}^{-1}$, and the reason for this is attributed to the interfering effect of both the soil texture and the levels of added magnesium and potassium in influencing the amount absorbed by the plant.




Figure (1): The effect of soil texture and the levels of magnesium and added potassium in the amount of potassium absorbed, mg K. plant^{-1} after 90 days of planting.

5. The effect of soil texture and levels of magnesium and added potassium in plant height, cm

Figure (2) shows the effect of soil texture, magnesium fertilizer, and potassium fertilizer in plant height in cm. The results showed that there were significant differences between the treatments. The results also indicated the superiority of the S_1 treatment due to its higher clay content ready-made potassium compared to S_2 , where the percentage of increase in plant height was In the S_1 treatment, 3.527% compared to the S_2 treatment, and this was confirmed

by (Hamid *et al.*, 2021) about the role of soil texture, the amount of clay and silt, and its mineral composition in the properties of the soil and its ability to supply plants. With nutrients and its effect in increasing plant height. As for the effect of magnesium fertilizer levels, the results showed that there were significant differences in plant height with an increase rate of 3.139, 6.688, 9.911 and 14.905% for the levels Mg₁, Mg₂, Mg₃ and Mg₄ compared to the control treatment Mg₀, respectively, and this is consistent with what was obtained (Mahmood *et al.*, 2020). who indicated that adding magnesium contributed to increasing plant height due to the role of magnesium in increasing the plant's photosynthesis process . The results also showed that adding potassium fertilizer at different levels led to a significant increase in plant height in the K₁₀₀ treatment compared to the control treatment K₀, by a rate of 6.891%. The increase in plant height can be attributed by increasing the levels of adding potassium fertilizer to the role of potassium in the accumulation of carbohydrates in the stem and increasing The number of nodes and their thickness, and thus the elongation of the stem, which is reflected positively in increasing the height of the plant, and this is consistent with the findings of (Al-Rawi & Al-Jumaili, 2018; Ali & Hamoud ,2022). As for the bilateral interaction between soil texture and magnesium levels, the results showed significant differences between the different treatments. The reason for this is attributed to the overlapping effect of soil texture and magnesium levels in plant height. The same trend also occurred with the bilateral interaction between the levels of magnesium and added potassium. The results showed significant differences between the different treatments. The reason for this is attributed to the interfering effect of the levels of magnesium and potassium fertilizer in plant height. As for the bilateral interaction between soil texture and potassium levels, the results showed significant differences between the different treatments. The reason for this is attributed to the overlapping effect of soil texture and potassium levels in plant height. As for the three-way interaction between soil texture and the levels of added magnesium and potassium, the results indicate that there are significant differences in plant height, as the highest rate of plant height in the S₁K₁₀₀ Mg₄ treatment reached 124.330 cm, while the lowest rate of soluble potassium in the S₂K₀Mg₀ treatment reached 98.330 cm. The reason for this is due to the interfering effect of soil texture and magnesium and potassium fertilizers on plant height.




Figure (2): The effect of soil texture and levels of magnesium and added potassium in plant height, cm after 90 days of planting.



6. Effect of soil texture and levels of magnesium and potassium added in leaf area cm^2

Shown in Figure (3) the effect of soil texture, magnesium fertilizer, and potassium fertilizer in leaf area cm^2 , the results showed that there were significant differences between the treatments. The results also indicated the superiority of the S_1 treatment due to its high clay content, which contributed to increasing the ready potassium compared to S_2 , where it reached The percentage increase in leaf area in the S_1 treatment was 21.044% compared to the S_2 treatment, and this was confirmed by (Hamid *et al.*, 2021) about the role of soil texture and the amount of clay and silt and its mineral composition in Soil characteristics and its ability to supply the plant with nutrients and its effect on increasing the leaf area of the plant, as there is a positive significant relationship with the amount of clay and silt and a negative relationship with sand in plant growth. As for the effect of magnesium fertilizer levels, the results showed that there were significant differences in the leaf area. The percentage increase reached 13.496, 26.660, 36.867 and 53.500% for the levels Mg_1 , Mg_2 , Mg_3 and Mg_4 compared to the control treatment Mg_0 , respectively. that The increase in the leaf area of the yellow corn plant is a result of increasing The level of magnesium addition is attributed to its effective role in vital processes within the plant, such as the process of photosynthesis and respiration, due to its inclusion in the chlorophyll molecule, in addition to its role in activating a number of enzymes, which led to an increase in the leaf area of the yellow corn plant. This is consistent with what was obtained by (Saaseea & Al-Amri, 2018). who indicated that the leaf area of yellow corn plants increased by increasing the level of magnesium addition. The results also showed that adding potassium fertilizer at different levels led to a significant increase in leaf area. The percentage increase in leaf area in the K_{100} treatment compared to the control treatment K_0 was 29.105%. The increase in leaf area can be attributed to the role of potassium in increasing growth rates. cells and improve vegetative growth, which reflects positively on the leaf area of the plant, and this is consistent with the findings of (Al-Rawi & Al-Jumaili, 2018). As for the bilateral interaction between soil texture and magnesium levels, the results showed significant differences between the different treatments. The reason for this is attributed to the overlapping effect of soil texture and magnesium levels in leaf area. The same trend also occurred with the bilateral interaction between the levels of added magnesium and potassium. The results showed significant differences between the different treatments in leaf area. The reason for this is attributed to the overlapping effect of the levels of magnesium and potassium fertilizer in increasing the amount of ready potassium and its release from the mineral layers, which contributed to its absorption by the plant . As for the bilateral interaction between soil texture and potassium levels, the results showed significant differences between the different treatments. The reason for this is attributed to the overlapping effect of soil texture and potassium levels in leaf area. As for the three-way interaction between soil texture and levels of added magnesium and potassium, the results indicate that there are significant differences in leaf area, as the highest average leaf area in the $S_1K_{100}Mg_4$ treatment reached 5684.00 cm^2 , while the lowest average leaf area in the $S_2K_0Mg_0$ treatment reached 2275.00 cm^2 . The reason for this is due to the interfering effect of soil texture and magnesium and potassium fertilizers.




Figure (3): Effect of soil texture and levels of added magnesium and potassium on leaf area cm^2 after 90 days of planting

CONCLUSION

The addition of magnesium affected the chemical equilibrium of Solube, exchangeable and non-exchangeable potassium due to its effect in the mineral composition.

REFERENCES

1. Al-Fadhly, J. T.M., Firas W. Ahmed & Abd, W. M. (2019). Effect of spraying with zinc and manganese at different stage of potato growth on quality of potato tuber (*Solanum tuberosum L.*) desiree class: as a role of friendly biochemical health. *Biochem Cell Arch*, 19 (2): 3955-3959.
2. Ali, W. M. & Hamoud, F. A.W. (2022). The effect of spraying organic emulsion (Appetizer) and nano-NPK with urea on some growth traits of three synthetic varieties of yellow corn. *Iraqi Journal of Market Research and Consumer Protection*, 14 (1): 108-117.
3. Al-Khalil, S. M.A., & Essa, S. K. (2020). Effect of sedimentation sources on the nature occurrence and distribution of the feldspar in some soil of alluvial plain Iraq. *Plant Archives*, 20 (2): 818-822.
4. AKhalil, S. M.A., & Essa, S. K. (2021). Effect of sedimentary source on the Properties of Sphericity and Roundness of feldspar minerals in some soils of the Alluvial Plain. *Indian Journal of Ecology*, 48 (1), 315-318.
5. Al-Marsoumi, A. E. & Jarallah, A. K. A. (2019). The effect of adding magnesium on the efficiency of phosphorus use in some phosphate fertilizers and in the absorption of magnesium and phosphorus in yellow corn (*Zea mays L.*). *Journal of Ethno-Agricultural Sciences*, 50 (5): 1302-1312.
6. Al-Obaidi, S. M.& Abdel Reda H. A. (2021). Evaluation of the combination of bacterial biofertilizer and vermicompost on the readiness of K, P, N and some growth standards of the bean plant *Phaseolus (vulgarisL.)*. *Iraqi Journal of Agricultural Sciences*, 52 (4), 960-970.
7. Al-Rawi, M. M. & Al-Jumaili, M. A.H. (2018). The effect of spraying potassium and zinc in the growth and yield of pods and okra seeds, *Iraqi Journal of Agricultural Sciences*, 49(6):1041-1048.
8. Al-Shamare, A. H.D. & Essa, S. K. (2020). The effect of sedimentation sources on the exchange properties of the clay particles of some soils in wasit and maysan governorates, *Plant Archives* 20 (2): 566-573



9. Al-Shamary, H. K., Al-Maamouri, A.B. D.S, & Sadeq, J. H . (2022). Effects of climatic variation on weathering intensity for the mineral composition in some Iraqi soils. *Caspian Journal of Environmental Sciences, Caspian Journal of environmental Sciences*. 20 (5): 991-1001.
10. Al- Taee, Z.A. & Al-Maamouri, AB. D.S. (2020). Study of the mineral compostion of some Iraqi Soils and its effect on zinc dsorption. *Plant Archives*, 20 (1): 769-776.
11. Aqrabi, H. S. Y. (2018). The effect of organic and mineral fertilizers on the readiness of potassium in the soil and the yield of chickpeas (*Cicer arietinum* L.). *Iraqi Journal of Agricultural Sciences*, 49 (2): 295-301.
12. Asaad, S. A. & Abdul Rasoul Q. J. (2017). An evaluation study of some methods for extracting ready soil potassium and its relationship to potassium absorbed from the barley crop. *Iraqi Journal of Agricultural Sciences*, 48 (6): 1697-1704.
13. Black, C.A. ED. (1965). Methods of soil analysis. part 2 chemical and microbiological properties. Am. Inc Madison. Wisconsin, USA. Soc. Agron.
14. Elfahdawi, W. A.T., Firas W. Ahmed & Cheyed, S. H. (2020). Effect of Agricultural Sulfur on Availability of NPK in Soil, growth and yield of corn (*Zea mays* L.). *Indian Journal of Ecology*, 47 (12): 200-205.
15. Hamid, S. M., Sahar, A. H., & Alaa, J. A. (2021). Physicochemical analyzes of selected soil samples from agricultural areas in Najaf Governorate. *Iraqi Journal of Market Research and Consumer Protection*, 13 (2):125-134.
16. Jackson, M. L. (1958). Soil chemical analysis. *Printice-Hall int. Englewood cliffs*.
17. Mahmood, Y. A., I. Q. Mohammed, Firas W. Ahmed & K. A. Wheib. 2020. Effect of organic, mineral fertilizers and foliar application of humic acid on growth and yield of corn (*Zea mays* L.). *Indian Journal of Ecology*. 47 Special Issue (10): 39-44.
18. Mahmood, Y. A., Firas W. Ahmed., Juma, S.S & Al-Arazah, W. M. (2019). Effect of solid and liquid organig fertilizer and spray with humic acid and nutrient uptake of nitrogen, phosphorus and potassium on growth, yield of cauliflower. *Plant Archives*, 19 (2): 1504-1509.
19. Nafawa, S. M. (2017). Study of the minerals controlling the solubility of calcium, magnesium, and iron and their relationship to soil management by adopting dissolution plans. *Iraqi Journal of Agricultural Sciences*, 48 (6):1715-1726.
20. Saaseea, K. K. & Al-Amri, N. J. K. (2018). The effect of spraying calcium and magnesium and fertilizing with humic acid on growth, yield and storage capacity of potato tubers. *Iraqi Journal of Agricultural Sciences*, 49 (5): 897-905.