



STUDY THE CORRELATION BETWEEN AFLATOXIN B1 AND MALE INFERTILITY IN IRAQI PATIENTS

Manar Talib Suhail ^{*1}, Muna T. Al-Musawi², Luma Hassan Alwan Al-obaigy³

¹Researcher, Biology department, college of sciences for women, university of Baghdad, Baghdad, Iraq.
mannar.taleb1102a@csu.uobaghdad.edu.iq

²Professor PhD., Institute of Genetic Engineering and Biotechnology for Post Graduate Studies, University of Baghdad, Baghdad, Iraq.
muna.t@csu.uobaghdad.edu.iq

³Assistant professor, Biology department, college of sciences for women, university of Baghdad, Baghdad, Iraq.
lumaha_bio@csu.uobaghdad.edu.iq

Received 26/ 5/ 2024, Accepted 10/ 10/ 2024, Published 30/ 6/ 2026

This work is licensed under a CCBY 4.0 <https://creativecommons.org/licenses/by/4.0>



ABASTRCT

The purpose of the investigation was to identify aflatoxin B1 (AFB1) in the blood serum and seminal fluid of 50 of patients from Iraqi male who suffering from infertility include (oligozoospermia, asthenozoospermia and teratozoospermia) with 16 fertile males as control depending on seminal fluid profile examination. The HPLC technique was used for AFB1 detection of the samples (serum and semen). The results showed that all the patient and control were gave positive detected of AFB1 50/50 (100%) and 16/16 (100%) respectively, in the serum at mean (81.08 ± 3.95) ng/ml at range (114-60) and (8.31 ± 0.63) ng/ml at range (5-12), respectively. Also detected AFB1 in the seminal fluid of patients at mean 39/50(78%) ng/ml at range (37-0), but not appear of AFB1 in any sample of control. There were highly significant differences ($P \leq 0.01$) between the mean level of AFB1 in the infertile male patients and control.

Key words: high performance liquid chromatography, infertility, oligozoospermia. mycotoxins.

دراسة العلاقة بين الأفلاتوكسين ب1 والعقم عند الذكور في المرضى العراقيين

منار طالب سهيل¹, منى تركي الموسوي², لمى علوان العبيدي³

¹باحث، قسم علوم الحياة، كلية العلوم للبنات، جامعة بغداد، بغداد، العراق mannar.taleb1102a@csu.uobaghdad.edu.iq

²استاذ دكتور، معهد الهندسة الوراثية والتقنيات الاحيائية للدراسات العليا، جامعة بغداد، بغداد، العراق muna.t@csu.uobaghdad.edu.iq

³استاذ مساعد دكتور، قسم علوم الحياة، كلية العلوم للبنات، جامعة بغداد، بغداد، العراق lumaha_bio@csu.uobaghdad.edu.iq

الخلاصة

هدفت الدراسة عن التحري على تشخيص سم الافلا ب 1 في مصل وسائل منوي لخمسون مريض من رجال عراقيين يعانون من عقم، شمل (قلة الحيامن، وهن الحيامن، تشوهات الحيامن) مع 16 رجل غير عقيم كسيطرة، بالاعتماد على فحص صورة السائل المنوي، استخدمت تقنية كروماتوغرافيا سائل عالي الدقة لتحري عن وجود سم الافلا ب 1 في جميع العينات (مصل وسائل منوي). أظهرت نتائج جميع المرضى والسيطرة تحري موجب لسم الافلا ب 1 50/50 (100%) و 16/16 (100%) في المصل على التوالي بمعدل (81.08 ± 3.95) جزء في البليون بمدى (114 - 60) ومعدل (8.31 ± 0.63) جزء في البليون بمدى (5 - 12)، على التوالي. كما تم التحري ايضا عن سم الافلا ب 1 في السائل المنوي للمرضى بمعدل 39/50 (78%) بمدى (37-0)، لكن لم يظهر سم الافلا ب 1 في أي عينة من السيطرة. وجد اختلاف احصائي عالي ($P \leq 0.01$) بين معدل مستوى سم الافلا ب 1 في الرجال العقيمين والسيطرة.

الكلمات المفتاحية: سموم فطرية 1، العقم، كروماتوغرافيا السائل عالي الدقة، قلة الحيامن.



INTRODUCTION

The inability to become pregnant after a year of consistent, unprotected sexual activity without the use of contraception is known as infertility (**Larsen, 2005**). "Male factor" infertility accounts for about 40–50% of male infertility, with 2% of cases having Sperm characteristics that are below ideal (**Kumar & singh, 2005**). Infertility has been associated with emotional, sociocultural, and physical difficulties (**Larsen, 2005**). Additionally, a high frequency of STDs, underdeveloped or nonexistent testes, and anomalies of the pituitary and hypothalamus are linked to infertility (**Agarwal et al. 2021**). In Iraq there were previous study to detect AFB1 in the serum and urine of Iraqi childhood and the result shows the high level of AFB1 in the serum and urine of children (**Al-musawi, 2020**).

Male infertility is most commonly caused by sperm morphology, movement issues, varicocele, difficulties ejaculating semen, low or nonexistent sperm counts, and genetic illnesses. Additionally, lifestyle decisions and environmental circumstances may have an effect (**Machen & Sandlow, 2020**). low motility of sperm (asthenozoospermia), reduced sperm quantity (oligozoospermia), reduced sperm vitality (necrozoospermia), aberrant sperm form (teratozoospermia), or any mix of these factors may be the cause (**Sharma, 2017**).

Iraq is one of countries that suffering from infertility, according to previous study in Iraq about the infertility rate, it has been estimated to be 3-4 out of 10 couples (35%) (**Al-Dujaily et al., 2017**). One of strong effect on infertility mycotoxins, mycotoxins can be defined as toxic secondary metabolites naturally produced by mold (*Aspergillus*, *fusarium* and *penicillium*) (**Turner & Snyder, 2021**). The fungi synthesize toxins after infesting crops which will be transmitted to the final food products. MTs effect on human healthy and causes problems, one of this problem infertility, Numerous studies have linked the rise in male infertility to the growing presence of these MTs in the environment. (**El. Khoury et al., 2019**). Studies shown that infertility rate has gotten worse from (42 to 48.5) million couples globally between 1990 and 2010, impacting 1 in 7 couples who are attempting to become pregnant. (**Stuppia et al., 2015**). The toxic category of fungal secondary metabolites known as AFs is naturally produced by *Aspergillus flavus* and *A. parasiticus*, with *A. nomius* producing them less frequently. AFs are formed at temperatures between 12 and 40 C and humidity levels between 3 and 18%. (**Arimboor, 2024**). Humans exposed to AFs run the risk of developing a number of diseases, including subacute and chronic consequences such cirrhosis, jaundice, chronic hepatitis, liver cancer, and hepatomegaly (**Hampikyan et al., 2010**) and immune system (**Jawad & Alwan, 2020**).

It has been reported that the AFB1 interferes with the enzymes and their substrate that are in charge of producing various hormones, hence impairing the activity of various endocrine glands. (**Gupta & Sharma, 2011**). AFs have the capacity to produce hormonal dysregulation that causes cellular toxicity, which has an immediate impact on reproduction (**Rossi et al., 2009**). According to earlier research, AFB1 disrupts the hypothalamo-pituitary testicular axis, which causes defective spermatozoa to be produced (**Supriya et al., 2014**).



In Iraq there are many study about detect MTs in human, Prior research has demonstrated that AFB1 in male human's causes delayed testicular development, testicular degeneration, reduced fertility, regressive morphological abnormalities in the testis, and dysfunction of the Leydig cell (Mohammed *et al.*, 2014). other study shows the presence of aflatoxins in serum and urine in Iraq (AL-Musawi, 2020) and other study detected aflatoxin in diet (Abu-Almaaly *et al.*, 2023; Muhyaddin *et al.*, 2015). Male infertility also effected by aflatoxins Also cause damage to the chromosome and gene by coupling to DNA, forming AFs-DNA complex lead to abnormalities in human sperm (Pribil, 2018).

Current study aimed to detect relationship between AFB1 and male infertility.

MATERIAL AND METHOD

Collection of seminal fluid samples

Freshly ejaculated samples of human sperm were obtained from males by masturbation and collected immediately into a clean, dry, and sterile disposable plastic Petri dish in a specially designated room. These samples from 50 patients and 16 healthy men according to seminal fluid examination and information acquired from every patient. The sample extracted from Kamal Al-Samarai Hospital in Baghdad and the period of sample collection from July 2022 to June 2023. The sample included infertile men who suffer from decrease in sperm count (oligozoospermi), sperm motile (asthenozoospermia), and abnormal sperm (teratozoospermia), while excluded from study; varicocele, azoospermia (no sperm production), viscosity samples and samples with pus cells. After 30-60 minutes, the samples were instantly delivered to the semen examination for each individual who had an acquaintance during 3-5 days of abstinence. Each semen sample was allowed to liquefy following WHO guidelines (Catanzariti *et al.*, 2013). Following full liquefaction, the sperm were examined macroscopically and microscopically in accordance with WHO standards from 1999 and 2010.

Collection of blood serum samples

Ten milliliters of patient's Venous blood were extracted from each patient while preserving sterility using a disposable syringe. After being put in a simple tube, the blood samples were left at room temperature for twenty minutes in order to allow them to coagulate. For 10 minutes at 3000 r.p.m., centrifugation was utilized to remove the serum from the clot. The resultant serum was divided in an Eppendorf tube and kept at -20°C until analysis to avoid repetitive cycles of freezing and thawing.

HPLC analysis

According to (McCoy *et al.*, 2008), the analysis was carried out in the labs of the Ministry of Science and Technology /Department of Water and Environment utilizing the HPLC technique for the detection and evaluation of mycotoxins AFB1 in serum and seminal fluid samples.

The reversed-phase HPLC (Cuknm, Germany type) on an Agilent 1100 system was linked in series with a fluorescence detector (366 nm excitation and 436 nm emission) that was



built around a diode-array UV detector at 362 nm. A C18 5 μm (150 $\times$ 4.6 mm) HPLC column (Waters Ltd., Watford, United Kingdom) was employed.

Chromatographic separation was obtained by a (5– 25%) ethanol linear gradient in water generated over a 25 min. period followed by isocratic elution with 25% ethanol in water, all at a flow rate of 1 ml/min.

The mobile phase was buffered with 5 mM triethylammonium formate (pH 3.0), and the column temperature was maintained at 35°C. After the eluted peaks were integrated, the standard curve was used to quantify AFB1. At 15.5 min, authentic AFB1 was eluted. The method's detection limit was five particles per milligram.

AFB1 Standard Curve

AFB1 was measured out to be about 0.5 mg, and the volume was added to a 100 ml volumetric flask until the concentration reached 5 ng/ml. To create a calibration curve, a number of concentrations were generated and injected into the HPLC using the dilution law ($C_1V_1=C_2V_2$).

To determine the retention period, standard AFB1 compounds were loaded into HPLC method devices at a concentration of 0.12 ng/ml (Fig. 1).




Fig. (1): Aflatoxin B1 standard curve.

STATISTICAL ANALYSIS

The tool SAS (**Statistical Analysis System, 2018**) was utilized to identify the impact of variation factors in the study parameters. The t-test was employed to compare means statistically. The chi-square test was used in this investigation to compare statistically significant percentages (0.05 and 0.01 likelihood).



RESULTS & DISSCUSSION

The present investigation is one of the few in Iraq and all world to detected the effect of aflatoxin b1 on human male fertility, there have been numerous published investigations on the identification of MTs in various food sources (Saadullah ,2018).

Detection of AFB1 in the serum

Table. (1) The results showed that all the patient and control were gave positive detection of AFB1 50/50 (100%) and {16/16 (100%)} respectively, in the serum at mean (81.08 $\pm$ 3.95) ng/ml at range (114-60) and (8.31 $\pm$ 0.63) ng/ml at range (5-12), respectively. by HPLC technique with mean 8.31 $\pm$ 0.63 ng/ml, there. were highly significantly differences * ($P \leq 0.01$). between this group, additionally, to maintain control.

Table (1): Distribution of AFB1 in the serum by HPLC technique.

Group	No of sample	Positive No. (%)	Mean by ng/ml	Range (min-mix)
Patients	50	50 (100%)	81.08 $\pm$ 3.95	60-114
Control	16	16 (100%)	8.31 $\pm$ 0.63	5-12
P-value	---	1.00 NS	0.0001 *	
* ($P \leq 0.01$).				

Detection of AFB1 in the seminal fluid

Table (2) show AFB1 in the seminal fluid of patients at mean 39/50(78%) ng/ml at range (37-0), but not appear of AFB1 in any sample of control. There were highly significantly differences ($P \leq 0.01$) between the mean level of AFB1 in the infertile male patients and control.

Table (2): Distribution of AFB1 in the seminal fluid by HPLC technique.

Group	No of sample	Positive No. (%)	Mean by ng/ml	Range (min-mix)
Patients	50	39 (78.00%)	15.7 $\pm$ 1.04	0-37
Control	16	0 (0.00%)	0.00 $\pm$ 0.00	0
P-value	---	0.0001 *	0.0001 *	
* ($P \leq 0.01$).				

This high prevalence of AFB1 in the semen of infertile group can be explained by the highest prevalence of aflatoxin B1 in different type of diet, the current observations conflict with a prior investigation by (Tang *et al.*, 2008) They found that, despite the paucity of research on the detection of AFB1 in human serum by HPLC technology, was the less sensitive



approach for detecting AFTs in human serum. Nevertheless, Hatem et al. (Hatem et al., 2005). Study by (Komsky-Elbaz, et al., 2018), that study exposure of seminal fluid to AFB1 in vitro, this studied show the direct on sperm viability by damage cell membrane lead to sperm dead and by effect on mitochondria lead to dysfunction apoptosis, calcium signaling, reactive oxygen species generation, and ATP synthesis. A physiological issue linked to changes in mitochondrial function includes infertility in both males and females (Komsky-Elbaz et al., 2018). Also effect on the DNA fragmentation lead male infertility (Ramalho-Santos et al., 2009). Other study on model revealed Boar seminal plasma was discovered to contain a significant concentration of AFB1 residue. AFB1 can also cause testicular lesions in mice, reducing the number and quality of their sperm, and ultimately affecting male fertility (Supriya et al., 2014). Exposure to AFB1 can lead to an imbalance between the oxidant and antioxidant enzyme systems, hence facilitating the excessive formation of reactive oxygen species (ROS) and consequent oxidative damage. (Kannan et al., 2014). Testicular lipid and DNA oxidative damage can disrupt spermatogenesis and the testis, which is linked to low-quality semen and increased infertility (Alahmar, 2019). Furthermore, oxidative stress in the testis has been widely shown to cause spermatogenic cells to undergo mitochondrial damage, apoptosis, and adverse alterations in sperm motility and concentration (Zhang et al., 2020).

CONCLUSION

Positive detection of AFB1 in the serum and seminal fluid of infertile male by using HPLC technique. This high prevalence of AFB1 in the semen of infertile group can be explained by the highest prevalence of AF B1 in different type of diet, presence of AFB1 in semen effect direct on male infertility on sperm count, sperm motility and morphology.

REFERENCES

1. Abu-Almaaly, R. A., Al-Musawi, A. T., & Mohammed, S. J. (2023). Estimation of some mycotoxins and heavy metals in the liver of important chickens available in the local markets. *Iraqi Journal of Market Research and Consumer Protection*, 15(2), 23-30,
2. Agarwal, A., Baskaran, S. Parekh. N, Cho. C-L, Henkel. R, Vij. S, Arafa. M, Panner Selva.MK and Shah, R. (2021). Male infertility. *Lancet* 397 319–333.
3. Alahmar, A. T. (2019). Role of oxidative stress in male infertility: an updated review. *Journal of human reproductive sciences*, 12(1), 4-18.
4. Al-Dujaily, S. S., Salah M. Al-Chalabi, N. Khadim, M. Abdul Mageed, and M. O. Abdul-Lateef. (2017). Effect of fertility blend administration on the oocytes quality and embryonic development using assisted reproductive technology in mice. *Journal of Biotechnology Research Center*, 11(1): 67-71.
5. Al-musawi. A. M. (2020). Detection of Aflatoxin B1 among Early and Middle Childhood Iraqi Patients. *Baghdad Science Journal*, 17(2) (SI), 0604-0604.
6. Arimboor, R. (2024). Metabolites and degradation pathways of microbial detoxification of aflatoxins: a review. *Mycotoxin Research*, 40(1), 71-83



7. Catanzariti, F., Cantoro, U., Lacetera, V., Muzzonigro, G., & Polito, M. (2013). Comparison between WHO (World Health Organization) 2010 and WHO 1999 parameters for semen analysis–interpretation of 529 consecutive samples. *Archivio Italiano di Urologia e Andrologia*, 85(3), 125-129.
8. El. Khoury, D., Fayjaloun, S., Nassar, M., Sahakian, J., & Aad, P. Y. (2019). Updates on the effect of mycotoxins on male reproductive efficiency in mammals. *Toxins*, 11(9), 515.
9. Francis, Sullibie, Kortei, Nii Korley, Sackey, Marian and Richard, Seidu A. (2024). "Aflatoxin B₁ induces infertility, fetal deformities, and potential therapies" *Open Medicine*, vol. 19, no. 1, pp. 20240907.
10. Gupta, R., & Sharma, V. (2011). Ameliorative effects of *Tinospora cordifolia* root extract on histopathological and biochemical changes induced by aflatoxin-B1 in mice kidney. *Toxicology international*, 18(2), 94.
11. Hatem, NL, Hassab, HM, Abd, AL- Rahman E, M.ELDeeb S A, and EL-Saeed A RL.(2005). Prevalence of afatoxins in blood and urin of gyptian infants with protein energy malnutrition. *Food and Nutr. Bull.*;26 (1): 49-56.
12. Hampikyan, H., Bingol, E. B., Cetin, O., & Colak, H. (2010). Determination of aflatoxin M1 levels in Turkish white, kashar and tulum cheeses. *Journal of Food, Agriculture and Environment*, 8, 13-15.
13. Jawad, B. J., & ALwan, M. J. (2020). Influence of mycotoxins on immune responses against *Salmonella typhimurium* infection of broiler chickens. *The Iraqi Journal of Agricultural Science*, 51(6), 1716-1725.
14. Kannan, K., Supriya, S., Adhithya, R., & Velazhahan, R. (2014). *Aspergillus flavus* and aflatoxin contamination of poultry feeds in Tamil Nadu, India. *International Journal of Agriculture, Environment and Biotechnology*, 7(2), 361-366.
15. Komsky-Elbaz, A., Saksier, M., & Roth, Z. (2018). Aflatoxin B1 impairs sperm quality and fertilization competence. *Toxicology*, 393, 42-50.
16. Kumar N and Singh AK. (2015). Trends of male factor infertility, an important cause of infertility: a review of literature. *Journal of Human Reproductive Sciences* 8 191–196.
17. Larsen U. (2005). Research on infertility: which definition should we use? *Fertility and Sterility* 83 846–852.
18. Machen, G. L., & Sandlow, J. I. (2020). Causes of male infertility. *Male Infertility: Contemporary Clinical Approaches, Andrology, ART and Antioxidants*, 3-14.
19. McCoy, L. F., Scholl, P. F., Sutcliffe, A. E., Kieszak, S. M., Powers, C. D., Rogers, H. S., ... & Schleicher, R. L. (2008). Human aflatoxin albumin adducts quantitatively compared by ELISA, HPLC with fluorescence detection, and HPLC with isotope dilution mass spectrometry. *Cancer Epidemiology Biomarkers & Prevention*, 17(7), 1653-1657.
20. Mohammed, M. N., Ameen, M. M., Mohammed, O. A., Al-Maghraby, O. M., Aziz, O. A., Ahmed, S. R., & Alsammani, M. A. (2014). The effect of aflatoxins on male reproduction. *Medical Archives*, 68(4), 272.



21. Muhyaddin, M., A. Al-Soufi, M., & T Al-Musawi, A. (2015). Determination of some Optimum Conditions for Bioremediation of Some Heavy Metals by *Saccharomyces cerevisiae*. *iraq journal of market research and consumer protection*, 7(2), 271-290.
22. Pribil, A. I. (2018). *Perils of the Fungal Kingdom: Mycotoxins in Food and Feed*. Louisiana State University and Agricultural & Mechanical College.
23. Ramalho-Santos, J., Varum, S., Amaral, S., Mota, P. C., Sousa, A. P., & Amaral, A. (2009). Mitochondrial functionality in reproduction: from gonads and gametes to embryos and embryonic stem cells. *Human reproduction update*, 15(5), 553-572.
24. SAS. (2018). *Statistical Analysis System, User's Guide*. Statistical. Version 9.6th ed. SAS. Inst. Inc. Cary. N.C. USA.
25. Saadullah. A. (2018). Mycoflora and incidence of Aflatoxin in wheat seeds from Duhok province, Kurdistan region of Iraq. *science journal of university of zakho*, 6 (.3), 78-81.
26. Sharma, A. (2017). Male infertility; evidences, risk factors, causes, diagnosis and management in human. *Annals of Clinical and Laboratory Research*, 5(3), 188.
27. Stuppia, L., Franzago, M., Ballerini, P., Gatta, V., & Antonucci, I. (2015). Epigenetics and male reproduction: the consequences of paternal lifestyle on fertility, embryo development, and children lifetime health. *Clinical epigenetics*, 7, 1-15.
28. Supriya, C., Girish, B. P., & Reddy, P. S. (2014). Aflatoxin B1-induced reproductive toxicity in male rats: possible mechanism of action. *International journal of toxicology*, 33(3), 155-161.
29. Tang, L., Tang, M., Xu, L., Luo, H., Huang, T., Yu, J., ... & Wang, J. S. (2008). Modulation of aflatoxin biomarkers in human blood and urine by green tea polyphenols intervention. *Carcinogenesis*, 29(2), 411-417.
30. Turner, P. C., & Snyder, J. A. (2021). Development and limitations of exposure biomarkers to dietary contaminants mycotoxins. *Toxins*, 13(5), 314.
31. Zhang, X., Cui, W., Wang, K. E., Chen, R., Chen, M., Lan, K., ... & Lan, X. (2020). Chlorpyrifos inhibits sperm maturation and induces a decrease in mouse male fertility. *Environmental Research*, 188, 109785.